


# The Online Math Open Fall Contest

November 4 – 15, 2016



# Acknowledgements

## Tournament Director

- James Lin

## Problem Authors

- Vincent Huang
- Yang Liu
- Michael Ren
- Ashwin Sah
- Tristan Shin
- Yannick Yao

## Website Manager

- Evan Chen
- Douglas Chen

## L<sup>A</sup>T<sub>E</sub>X/Python Geek

- Evan Chen

# Contest Information

These rules supersede any rules found elsewhere about the OMO. Please send any further questions directly to the OMO Team at [OnlineMathOpenTeam@gmail.com](mailto:OnlineMathOpenTeam@gmail.com).

## Team Registration and Eligibility

Students may compete in teams of up to four people, but no student can belong to more than one team. Participants must not have graduated from high school (or the equivalent secondary school institution in other countries). Teams need not remain the same between the Fall and Spring contests, and students are permitted to participate in whichever contests they like.

**Only one member on each team needs to register an account on the website.** Please check the website, [http://internetolympiad.org/pages/14-omo\\_info](http://internetolympiad.org/pages/14-omo_info), for registration instructions.

*Note:* when we say “up to four”, we really do mean “up to”! Because the time limit is so long, partial teams are not significantly disadvantaged, and we welcome their participation.

## Contest Format and Rules

The 2016 Fall Contest will consist of 30 problems; the answer to each problem will be an integer between 0 and  $2^{31} - 1 = 2147483647$  inclusive. The contest window will be

November 4 – 15, 2016

from 7PM ET on the start day to **7PM ET on the end day**. There is no time limit other than the contest window.

1. Four-function calculators (calculators which can perform only the four basic arithmetic operations) are permitted on the Online Math Open. **Any other computational aids, including scientific calculators, graphing calculators, or computer programs is prohibited.** All problems on the Online Math Open are solvable without a calculator. Four-function calculators are permitted only to help participants reduce computation errors.
2. Drawing aids such as graph paper, ruler, and compass are permitted. However, electronic drawing aids, such as Geogebra and graphing calculators, are not allowed. Print and electronic publications are also not allowed.
3. Members of different teams cannot communicate with each other about the contest while the contest is running.
4. Your score is the number of questions answered correctly; that is, every problem is worth one point. Ties will be broken based on the “hardest” problem that a team answered correctly. Remaining ties will be broken by the second hardest problem solved, and so on. (Problem  $m$  is harder than problem  $n$  if fewer teams solve problem  $m$  OR if the number of solves is equal and  $m > n$ .)
5. *Participation in the Online Math Open is free.*

## Clarifications and Results

Clarifications will be posted as they are answered. For the most recent contests, they will be posted at [http://internetolympiad.org/pages/n/omo\\_problems](http://internetolympiad.org/pages/n/omo_problems). If you have a question about problem wording, please email [OnlineMathOpenTeam@gmail.com](mailto:OnlineMathOpenTeam@gmail.com) with “Clarification” in the subject. We have the right to deny clarification requests that we feel we cannot answer.

After the contest is over, we will release the answers to the problems within the next day. Please do not discuss the test until answers are released. If you have a protest about an answer, you may send an email to [OnlineMathOpenTeam@gmail.com](mailto:OnlineMathOpenTeam@gmail.com). (Include “Protest” in the subject). Results will be released in the following weeks. (Results will be counted only for teams that submit answers at least once. Teams that only register an account will not be listed in the final rankings.)

**OMO Fall 2016**  
**November 4 – 15, 2016**

1. Kevin is in first grade, so his teacher asks him to calculate  $20 + 1 \cdot 6 + k$ , where  $k$  is a real number revealed to Kevin. However, since Kevin is rude to his Aunt Sally, he instead calculates  $(20+1) \cdot (6+k)$ . Surprisingly, Kevin gets the correct answer! Assuming Kevin did his computations correctly, what was his answer?
2. Yang has a standard 6-sided die, a standard 8-sided die, and a standard 10-sided die. He tosses these three dice simultaneously. The probability that the three numbers that show up form the side lengths of a right triangle can be expressed as  $\frac{m}{n}$ , for relatively prime positive integers  $m$  and  $n$ . Find  $100m + n$ .
3. In a rectangle  $ABCD$ , let  $M$  and  $N$  be the midpoints of sides  $BC$  and  $CD$ , respectively, such that  $AM$  is perpendicular to  $MN$ . Given that the length of  $AN$  is 60, the area of rectangle  $ABCD$  is  $m\sqrt{n}$  for positive integers  $m$  and  $n$  such that  $n$  is not divisible by the square of any prime. Compute  $100m + n$ .

4. Let  $G = 10^{10^{100}}$  (a.k.a. a googolplex). Then

$$\log_{(\log_{(\log_{10} G)} G)} G$$

can be expressed in the form  $\frac{m}{n}$  for relatively prime positive integers  $m$  and  $n$ . Determine the sum of the digits of  $m + n$ .

5. Jay notices that there are  $n$  primes that form an arithmetic sequence with common difference 12. What is the maximum possible value for  $n$ ?
6. For a positive integer  $n$ , define  $n? = 1^n \cdot 2^{n-1} \cdot 3^{n-2} \cdots (n-1)^2 \cdot n^1$ . Find the positive integer  $k$  for which  $7?9? = 5?k?$ .
7. The 2016 players in the Gensokyo Tennis Club are playing Up and Down the River. The players first randomly form 1008 pairs, and each pair is assigned to a tennis court (The courts are numbered from 1 to 1008). Every day, the two players on the same court play a match against each other to determine a winner and a loser. For  $2 \leq i \leq 1008$ , the winner on court  $i$  will move to court  $i - 1$  the next day (and the winner on court 1 does not move). Likewise, for  $1 \leq j \leq 1007$ , the loser on court  $j$  will move to court  $j + 1$  the next day (and the loser on court 1008 does not move). On Day 1, Reimu is playing on court 123 and Marisa is playing on court 876. Find the smallest positive integer value of  $n$  for which it is possible that Reimu and Marisa play one another on Day  $n$ .
8. For a positive integer  $n$ , define the  $n$ th triangular number  $T_n$  to be  $\frac{n(n+1)}{2}$ , and define the  $n$ th square number  $S_n$  to be  $n^2$ . Find the value of

$$\sqrt{S_{62} + T_{63} \sqrt{S_{61} + T_{62} \sqrt{\cdots \sqrt{S_2 + T_3 \sqrt{S_1 + T_2}}}}}$$

9. In quadrilateral  $ABCD$ ,  $AB = 7$ ,  $BC = 24$ ,  $CD = 15$ ,  $DA = 20$ , and  $AC = 25$ . Let segments  $AC$  and  $BD$  intersect at  $E$ . What is the length of  $EC$ ?
10. Let  $a_1 < a_2 < a_3 < a_4$  be positive integers such that the following conditions hold:
  - $\gcd(a_i, a_j) > 1$  holds for all integers  $1 \leq i < j \leq 4$ .
  - $\gcd(a_i, a_j, a_k) = 1$  holds for all integers  $1 \leq i < j < k \leq 4$ .

Find the smallest possible value of  $a_4$ .

11. Let  $f$  be a random permutation on  $\{1, 2, \dots, 100\}$  satisfying  $f(1) > f(4)$  and  $f(9) > f(16)$ . The probability that  $f(1) > f(16) > f(25)$  can be written as  $\frac{m}{n}$  where  $m$  and  $n$  are relatively prime positive integers. Compute  $100m + n$ .

Note: In other words,  $f$  is a function such that  $\{f(1), f(2), \dots, f(100)\}$  is a permutation of  $\{1, 2, \dots, 100\}$ .

**OMO Fall 2016**  
**November 4 – 15, 2016**

12. For each positive integer  $n \geq 2$ , define  $k(n)$  to be the largest integer  $m$  such that  $(n!)^m$  divides 2016!. What is the minimum possible value of  $n + k(n)$ ?
13. Let  $A_1B_1C_1$  be a triangle with  $A_1B_1 = 16$ ,  $B_1C_1 = 14$ , and  $C_1A_1 = 10$ . Given a positive integer  $i$  and a triangle  $A_iB_iC_i$  with circumcenter  $O_i$ , define triangle  $A_{i+1}B_{i+1}C_{i+1}$  in the following way:
- $A_{i+1}$  is on side  $B_iC_i$  such that  $C_iA_{i+1} = 2B_iA_{i+1}$ .
  - $B_{i+1} \neq C_i$  is the intersection of line  $A_iC_i$  with the circumcircle of  $O_iA_{i+1}C_i$ .
  - $C_{i+1} \neq B_i$  is the intersection of line  $A_iB_i$  with the circumcircle of  $O_iA_{i+1}B_i$ .

Find

$$\left( \sum_{i=1}^{\infty} [A_iB_iC_i] \right)^2.$$

Note:  $[K]$  denotes the area of  $K$ .

14. In Yang's number theory class, Michael K, Michael M, and Michael R take a series of tests. Afterwards, Yang makes the following observations about the test scores:
- Michael K had an average test score of 90, Michael M had an average test score of 91, and Michael R had an average test score of 92.
  - Michael K took more tests than Michael M, who in turn took more tests than Michael R.
  - Michael M got a higher total test score than Michael R, who in turn got a higher total test score than Michael K. (The total test score is the sum of the test scores over all tests)

What is the least number of tests that Michael K, Michael M, and Michael R could have taken combined?

15. Two bored millionaires, Bilion and Trilion, decide to play a game. They each have a sufficient supply of \$1, \$2, \$5, and \$10 bills. Starting with Bilion, they take turns putting one of the bills they have into a pile. The game ends when the bills in the pile total exactly \$1,000,000, and whoever makes the last move wins the \$1,000,000 in the pile (if the pile is worth more than \$1,000,000 after a move, then the person who made the last move loses instead, and the other person wins the amount of cash in the pile). Assuming optimal play, how many dollars will the winning player gain?
16. For her zeroth project at Magic School, Emilia needs to grow six perfectly-shaped apple trees. First she plants six tree saplings at the end of Day 0. On each day afterwards, Emilia attempts to use her magic to turn each sapling into a perfectly-shaped apple tree, and for each sapling she succeeds in turning it into a perfectly-shaped apple tree that day with a probability of  $\frac{1}{2}$ . (Once a sapling is turned into a perfectly-shaped apple tree, it will stay a perfectly-shaped apple tree.) The expected number of days it will take Emilia to obtain six perfectly-shaped apple trees is  $\frac{m}{n}$  for relatively prime positive integers  $m$  and  $n$ . Find  $100m + n$ .
17. Let  $n$  be a positive integer.  $S$  is a set of points such that the points in  $S$  are arranged in a regular 2016-simplex grid, with an edge of the simplex having  $n$  points in  $S$ . (For example, the 2-dimensional analog would have  $\frac{n(n+1)}{2}$  points arranged in an equilateral triangle grid). Each point in  $S$  is labeled with a real number such that the following conditions hold:
- Not all the points in  $S$  are labeled with 0.
  - If  $\ell$  is a line that is parallel to an edge of the simplex and that passes through at least one point in  $S$ , then the labels of all the points in  $S$  that are on  $\ell$  add to 0.
  - The labels of the points in  $S$  are symmetric along any such line  $\ell$ .

**OMO Fall 2016**  
**November 4 – 15, 2016**

Find the smallest positive integer  $n$  such that this is possible.

Note: A regular 2016-simplex has 2017 vertices in 2016-dimensional space such that the distances between every pair of vertices are equal.

18. Find the smallest positive integer  $k$  such that there exist positive integers  $M, O > 1$  satisfying

$$(O \cdot M \cdot O)^k = (O \cdot M) \cdot \underbrace{(N \cdot O \cdot M) \cdot (N \cdot O \cdot M) \cdots (N \cdot O \cdot M)}_{2016 \text{ } (N \cdot O \cdot M)\text{s}},$$

where  $N = O^M$ .

Note: This is edited from the previous text, which did not clarify that  $NOM$  represented  $N \cdot O \cdot M$ , for example.

19. Let  $S$  be the set of all polynomials  $Q(x, y, z)$  with coefficients in  $\{0, 1\}$  such that there exists a homogeneous polynomial  $P(x, y, z)$  of degree 2016 with integer coefficients and a polynomial  $R(x, y, z)$  with integer coefficients so that

$$P(x, y, z)Q(x, y, z) = P(yz, zx, xy) + 2R(x, y, z)$$

and  $P(1, 1, 1)$  is odd. Determine the size of  $S$ .

Note: A homogeneous polynomial of degree  $d$  consists solely of terms of degree  $d$ .

20. For a positive integer  $k$ , define the sequence  $\{a_n\}_{n \geq 0}$  such that  $a_0 = 1$  and for all positive integers  $n$ ,  $a_n$  is the smallest positive integer greater than  $a_{n-1}$  for which  $a_n \equiv ka_{n-1} \pmod{2017}$ . What is the number of positive integers  $1 \leq k \leq 2016$  for which  $a_{2016} = 1 + \binom{2017}{2}$ ?
21. Mark the Martian and Bark the Bartian live on planet Blok, in the year 2019. Mark and Bark decide to play a game on a  $10 \times 10$  grid of cells. First, Mark randomly generates a subset  $S$  of  $\{1, 2, \dots, 2019\}$  with  $|S| = 100$ . Then, Bark writes each of the 100 integers in a different cell of the  $10 \times 10$  grid. Afterwards, Bark constructs a solid out of this grid in the following way: for each grid cell, if the number written on it is  $n$ , then she stacks  $n$   $1 \times 1 \times 1$  blocks on top of one other in that cell. Let  $B$  be the largest possible surface area of the resulting solid, including the bottom of the solid, over all possible ways Bark could have inserted the 100 integers into the grid of cells. Find the expected value of  $B$  over all possible sets  $S$  Mark could have generated.
22. Let  $ABC$  be a triangle with  $AB = 3$  and  $AC = 4$ . It is given that there does not exist a point  $D$ , different from  $A$  and not lying on line  $BC$ , such that the Euler line of  $ABC$  coincides with the Euler line of  $DBC$ . The square of the product of all possible lengths of  $BC$  can be expressed in the form  $m + n\sqrt{p}$ , where  $m$ ,  $n$ , and  $p$  are positive integers and  $p$  is not divisible by the square of any prime. Find  $100m + 10n + p$ .

Note: For this problem, consider every line passing through the center of an equilateral triangle to be an Euler line of the equilateral triangle. Hence, if  $D$  is chosen such that  $DBC$  is an equilateral triangle and the Euler line of  $ABC$  passes through the center of  $DBC$ , then consider the Euler line of  $ABC$  to coincide with "the" Euler line of  $DBC$ .

23. Let  $\mathbb{N}$  denote the set of positive integers. Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be a function such that the following conditions hold:

- For any  $n \in \mathbb{N}$ , we have  $f(n) | n^{2016}$ .
- For any  $a, b, c \in \mathbb{N}$  satisfying  $a^2 + b^2 = c^2$ , we have  $f(a)f(b) = f(c)$ .

Over all possible functions  $f$ , determine the number of distinct values that can be achieved by  $f(2014) + f(2) - f(2016)$ .

**OMO Fall 2016**  
**November 4 – 15, 2016**

24. Let  $P(x, y)$  be a polynomial such that  $\deg_x(P), \deg_y(P) \leq 2020$  and

$$P(i, j) = \binom{i+j}{i}$$

over all  $2021^2$  ordered pairs  $(i, j)$  with  $0 \leq i, j \leq 2020$ . Find the remainder when  $P(4040, 4040)$  is divided by 2017.

Note:  $\deg_x(P)$  is the highest exponent of  $x$  in a nonzero term of  $P(x, y)$ .  $\deg_y(P)$  is defined similarly.

25. Let  $X_1X_2X_3$  be a triangle with  $X_1X_2 = 4, X_2X_3 = 5, X_3X_1 = 7$ , and centroid  $G$ . For all integers  $n \geq 3$ , define the set  $S_n$  to be the set of  $n^2$  ordered pairs  $(i, j)$  such that  $1 \leq i \leq n$  and  $1 \leq j \leq n$ . Then, for each integer  $n \geq 3$ , when given the points  $X_1, X_2, \dots, X_n$ , randomly choose an element  $(i, j) \in S_n$  and define  $X_{n+1}$  to be the midpoint of  $X_i$  and  $X_j$ . The value of

$$\sum_{i=0}^{\infty} \left( \mathbb{E} [X_{i+4} G^2] \left( \frac{3}{4} \right)^i \right)$$

can be expressed in the form  $p + q \ln 2 + r \ln 3$  for rational numbers  $p, q, r$ . Let  $|p| + |q| + |r| = \frac{m}{n}$  for relatively prime positive integers  $m$  and  $n$ . Compute  $100m + n$ .

Note:  $\mathbb{E}(x)$  denotes the expected value of  $x$ .

26. Let  $ABC$  be a triangle with  $BC = 9, CA = 8$ , and  $AB = 10$ . Let the incenter and incircle of  $ABC$  be  $I$  and  $\gamma$ , respectively, and let  $N$  be the midpoint of major arc  $BC$  of the circumcircle of  $ABC$ . Line  $NI$  meets the circumcircle of  $ABC$  a second time at  $P$ . Let the line through  $I$  perpendicular to  $AI$  meet segments  $AB, AC$ , and  $AP$  at  $C_1, B_1$ , and  $Q$ , respectively. Let  $B_2$  lie on segment  $CQ$  such that line  $B_1B_2$  is tangent to  $\gamma$ , and let  $C_2$  lie on segment  $BQ$  such that line  $C_1C_2$  tangent to  $\gamma$ . The length of  $B_2C_2$  can be expressed in the form  $\frac{m}{n}$  for relatively prime positive integers  $m$  and  $n$ . Determine  $100m + n$ .
27. Compute the number of monic polynomials  $q(x)$  with integer coefficients of degree 12 such that there exists an integer polynomial  $p(x)$  satisfying  $q(x)p(x) = q(x^2)$ .
28. Let  $ABC$  be a triangle with  $AB = 34, BC = 25$ , and  $CA = 39$ . Let  $O, H$ , and  $\omega$  be the circumcenter, orthocenter, and circumcircle of  $\triangle ABC$ , respectively. Let line  $AH$  meet  $\omega$  a second time at  $A_1$  and let the reflection of  $H$  over the perpendicular bisector of  $BC$  be  $H_1$ . Suppose the line through  $O$  perpendicular to  $A_1O$  meets  $\omega$  at two points  $Q$  and  $R$  with  $Q$  on minor arc  $AC$  and  $R$  on minor arc  $AB$ . Denote  $\mathcal{H}$  as the hyperbola passing through  $A, B, C, H, H_1$ , and suppose  $HO$  meets  $\mathcal{H}$  again at  $P$ . Let  $X, Y$  be points with  $XH \parallel AR \parallel YP, XP \parallel AQ \parallel YH$ . Let  $P_1, P_2$  be points on the tangent to  $\mathcal{H}$  at  $P$  with  $XP_1 \parallel OH \parallel YP_2$  and let  $P_3, P_4$  be points on the tangent to  $\mathcal{H}$  at  $H$  with  $XP_3 \parallel OH \parallel YP_4$ . If  $P_1P_4$  and  $P_2P_3$  meet at  $N$ , and  $ON$  may be written in the form  $\frac{a}{b}$  where  $a, b$  are positive coprime integers, find  $100a + b$ .
29. Let  $n$  be a positive integer. Yang the Saltant Sanguivorous Shearling is on the side of a very steep mountain that is embedded in the coordinate plane. There is a blood river along the line  $y = x$ , which Yang may reach but is not permitted to go above (i.e. Yang is allowed to be located at  $(2016, 2015)$  and  $(2016, 2016)$ , but not at  $(2016, 2017)$ ). Yang is currently located at  $(0, 0)$  and wishes to reach  $(n, 0)$ . Yang is permitted only to make the following moves:
- Yang may *spring*, which consists of going from a point  $(x, y)$  to the point  $(x, y + 1)$ .
  - Yang may *stroll*, which consists of going from a point  $(x, y)$  to the point  $(x + 1, y)$ .
  - Yang may *sink*, which consists of going from a point  $(x, y)$  to the point  $(x, y - 1)$ .

**OMO Fall 2016**  
**November 4 – 15, 2016**

In addition, whenever Yang does a *sink*, he breaks his tiny little legs and may no longer do a *spring* at any time afterwards. Yang also expends a lot of energy doing a *spring* and gets bloodthirsty, so he must visit the blood river at least once afterwards to quench his bloodthirst. (So Yang may still *spring* while bloodthirsty, but he may not finish his journey while bloodthirsty.) Let there be  $a_n$  different ways for which Yang can reach  $(n, 0)$ , given that Yang is permitted to pass by  $(n, 0)$  in the middle of his journey. Find the 2016th smallest positive integer  $n$  for which  $a_n \equiv 1 \pmod{5}$ .

30. Let  $P_1(x), P_2(x), \dots, P_n(x)$  be monic, non-constant polynomials with integer coefficients and let  $Q(x)$  be a polynomial with integer coefficients such that

$$x^{2^{2016}} + x + 1 = P_1(x)P_2(x) \dots P_n(x) + 2Q(x).$$

Suppose that the maximum possible value of  $2016n$  can be written in the form  $2^{b_1} + 2^{b_2} + \dots + 2^{b_k}$  for nonnegative integers  $b_1 < b_2 < \dots < b_k$ . Find the value of  $b_1 + b_2 + \dots + b_k$ .